

# Computer Science Department

## TECHNICAL REPORT

Subsequently published in Proc. Amer. Math. Soc., 100  
(1987), 319-321. MR 89b:47013a

AN INEQUALITY FOR SELF-ADJOINT OPERATORS  
ON A HILBERT SPACE

By

Herbert J. Bernstein  
December 1985

Technical Report #190

### NEW YORK UNIVERSITY



Department of Computer Science  
Courant Institute of Mathematical Sciences  
251 MERCER STREET, NEW YORK, N.Y. 10012

**AN INEQUALITY FOR SELF-ADJOINT OPERATORS  
ON A HILBERT SPACE**

**By**

**Herbert J. Bernstein**

**December 1985**

**Technical Report #190**

# An Inequality for Self-Adjoint Operators on a Hilbert Space

by Herbert J. Bernstein<sup>†</sup>

## ABSTRACT

An elementary inequality of use in testing convergence of eigenvector calculations is proven. If  $e_\lambda$  is a unit eigenvector corresponding to eigenvalue  $\lambda$  of self-adjoint operator  $A$  on a Hilbert space  $H$ , then

$$|(g, e_\lambda)|^2 \leq \frac{\|g\|^2 \|Ag\|^2 - (g, Ag)^2}{\|(A - \lambda I)g\|^2}$$

for all  $g$  in  $H$  for which  $Ag \neq \lambda g$ . Equality holds only when the component of  $g$  orthogonal to  $e_\lambda$  is also an eigenvector of  $A$ .

When computing eigenvectors of real symmetric matrices by iterative techniques, convergence is usually assumed on the basis of stagnation of Rayleigh quotients. (e.g. see [2]). In most cases this is satisfactory. However, when dealing with pathologically close eigenvalues, significant components orthogonal to the desired vector may remain [3, p 174]. In such cases one may try to estimate the size of those orthogonal components to decide if, say, Richardson's purification is needed. In this paper we present an estimator which has been of value in such calculations. The proof is elementary and the author suspects that it is not new, but is able to find no prior publication nor use within standard eigenvector programs.

Before proving the inequality, we state a lemma which can be derived directly from the equations used in proving Cauchy's or Schwartz's inequality. [1, p. 16]. We include an abstract proof here for completeness.

**Lemma.** Let  $H$  be a Hilbert space. Let  $A$  be a self-adjoint operator on  $H$ . Let  $x$  be a vector in  $H$ . Let  $\tau$  be real, then

1980 *Mathematics Subject Classification*. Primary 65F15, Secondary 65J10.

<sup>†</sup>Research partially supported by the Department of Energy under contract DEACO276ERO3077-V, the National Science Foundation, under contract DCR-8511302, and the Office of Naval Research, under contract N00014-82-K-0381.

$$\|x\|^2 \|Ax\|^2 - (x, Ax)^2 = \|x\|^2 \|(A - \tau I)x\|^2 - (x, (A - \tau I)x)^2$$

**Proof.**

$$\begin{aligned} & \|x\|^2 \|(A - \tau I)x\|^2 - (x, (A - \tau I)x)^2 \\ &= \|x\|^2 (\|Ax\|^2 - 2\tau(x, Ax) + \tau^2 \|x\|^2) \\ &\quad - ((x, Ax)^2 - 2\tau(x, Ax) \|x\|^2 + \tau^2 \|x\|^4) \\ &= \|x\|^2 \|Ax\|^2 - 2\tau(x, Ax) \|x\|^2 + \tau^2 \|x\|^4 \\ &\quad - (x, Ax)^2 + 2\tau(x, Ax) \|x\|^2 - \tau^2 \|x\|^4 \\ &= \|x\|^2 \|Ax\|^2 - (x, Ax)^2 \end{aligned}$$

**Q.E.D.**

Now we can prove the desired inequality.

**Theorem 1.** Let  $H$  be a Hilbert space. Let  $A$  be a self-adjoint operator on  $H$ . Let  $e_\lambda$  be a unit eigenvector of  $A$  with corresponding eigenvalue  $\lambda$ . Let  $f$  be orthogonal to  $A$ , then for any  $g = \alpha e_\lambda + f$  with  $\alpha$  real, either  $Ag = \lambda g$ , or

$$\alpha^2 \leq \frac{\|g\|^2 \|Ag\|^2 - (g, Ag)^2}{\|(A - \lambda I)g\|^2}$$

**Proof.** By the lemma, we may replace  $A$  in the numerator by  $A - \lambda I$ , allowing us to assume  $\lambda = 0$  without loss of generality. However, in that case

$$Ag = Af$$

$$(g, Ag) = (f, Af)$$

and

$$\|Ag\| = \|Af\|$$

so that

$$\begin{aligned} & \frac{\|g\|^2 \|Ag\|^2 - (g, Ag)^2}{\|Ag\|^2} \\ &= \frac{(\alpha^2 + \|f\|^2) \|Af\|^2 - (f, Af)^2}{\|Af\|^2} \\ &= \alpha^2 + \frac{\|f\|^2 \|Af\|^2 - (f, Af)^2}{\|Af\|^2} \end{aligned}$$

$$\geq \alpha^2$$

by the Schwartz inequality. *Q.E.D.*

Notice that the inequality is strict unless the Schwartz inequality on  $(f, Af)$  is an equality, *i.e.* when  $f$  is an eigenvector of  $A$ . It then follows immediately that

**Corollary.** If  $e_\lambda$  is a unit eigenvector corresponding to eigenvalue  $\lambda$  of self-adjoint operator  $A$  on a Hilbert space  $H$  then

$$|(g, e_\lambda)|^2 \leq \frac{\|g\|^2 \|Ag\|^2 - (g, Ag)^2}{\|(A - \lambda I)g\|^2}$$

for all  $g$  in  $H$  for which  $Ag \neq \lambda g$ . Equality will hold if and only if the component of  $g$  orthogonal to  $e_\lambda$  is also an eigenvector of  $A$ .

We now show that the bound given is actually the best of the bounds that could be obtained by taking quotients of a class of shifted inner products.

**Theorem 2.** Under the hypotheses of theorem 1, let  $\tau$  be real, then either  $Ag = \lambda g$  or

$$\frac{\|g\|^2 \|Ag\|^2 - (g, Ag)^2}{\|(A - \lambda I)g\|^2} \leq \frac{\|(A - \tau I)g\|^2}{(\lambda - \tau)^2}$$

**Proof.** Since  $\frac{a}{b} \leq \frac{c}{d}$ , for  $b, d > 0$  if and only if  $bc - ad \geq 0$ , we define  $B = A - \lambda I$  and, using the lemma, compute

$$\begin{aligned} & \|(A - \lambda I)g\|^2 \|(A - \tau I)g\|^2 - (\lambda - \tau)^2 (\|g\|^2 \|Ag\|^2 - (g, Ag)^2) \\ &= \|Bg\|^2 \|(B + (\lambda - \tau)I)g\|^2 - (\lambda - \tau)^2 (\|g\|^2 \|Bg\|^2 - (g, Bg)^2) \\ &= \|Bg\|^4 + 2(\lambda - \tau) \|Bg\|^2 + (\lambda - \tau)^2 \|g\|^2 \|Bg\|^2 \\ &\quad - (\lambda - \tau)^2 \|g\|^2 \|Bg\|^2 + (\lambda - \tau)^2 (g, Bg)^2 \\ &= (\|Bg\|^2 + (\lambda - \tau)(g, Bg))^2 \geq 0 \end{aligned}$$

with equality only when  $(g, (A - \lambda I)g) \neq 0$  and

$$\tau = \lambda + \frac{\|(A - \lambda I)g\|^2}{(g, (A - \lambda I)g)}$$

***Q.E.D.***

## References

1. G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, England, 1934.
2. H. Rutishauser, "Simultaneous Iteration Method for Symmetric Matrices," *Numer. Math.*, vol. 16, pp. 205-223, 1970.
3. J. H. Wilkinson, *The Algebraic Eigenvalue Problem*, Clarendon Press, Oxford, England, 1965.

Courant Institute of Mathematical Sciences, New York University, 251 Mercer Street, New York, New York 10012